

This question paper contains 3 printed pages]

**NA—41—2023**

**FACULTY OF SCIENCE**

**B.Sc. (Third Year) (Sixth Semester) EXAMINATION**

**NOVEMBER/DECEMBER, 2023**

**(CBCS/New Pattern)**

**MATHEMATICS**

**Paper XVI**

**(Integral Transforms)**

**(Monday, 11-12-2023)**

**Time : 10.00 a.m. to 12.00 noon**

*Time—Two Hours*

*Maximum Marks—40*

*N.B. :— (i) All questions are compulsory.*

*(ii) Figures to the right indicate full marks.*

1. If  $L[f(t)]$  denote the Laplace transform of  $f(t)$ , then prove that : 15

$$L[f^n(t)] = s^n L[f(t)] - s^{n-1} f(0) - s^{n-2} f'(0) - s^{n-3} f''(0) \dots - f^{n-1}(0).$$

Hence find the Laplace transform of  $t^2 \cos at$ .

*Or*

(a) Find the inverse Laplace transform of : 8

$$\frac{s-1}{s^2-6s+25}$$

P.T.O.

WT

( 2 )

NA—41—2023

(b) Find the inverse Laplace transform of :

7

$$\frac{s^2}{(s^2 + a^2)(s^2 + b^2)}.$$

2. Prove the Fourier sine and cosine integrals :

15

$$f(x) = \frac{2}{\pi} \int_0^{\infty} \sin ux \, du \int_0^{\infty} f(t) \cdot \sin ut \, dt$$

and

$$f(x) = \frac{2}{\pi} \int_0^{\infty} \cos ux \, du \int_0^{\infty} f(t) \cos ut \, dt .$$

Or

(a) Using the Laplace transforms, find the solution of the initial value problem :

8

$$y'' + 25y = 10 \cos 5t ,$$

$$y(0) = 2 , \quad y'(0) = 0 .$$

(b) Solve the initial value problem :

7

$$2y'' + 5y' + 2y = e^{-2t} ,$$

$$y(0) = 1 , \quad y'(0) = 1 .$$

WT

( 3 )

NA—41—2023

3. Attempt any *two* of the following :

5 marks each

(a) Find the Laplace transform of  $\cos^2 t$ .

(b) Find the inverse Laplace transform of :

$$\frac{s^2 + 3}{s(s^2 + 9)}.$$

(c) Applying convolution, solve the following initial value problem :

$$y'' + y = \sin 3t,$$

$$y(0) = 0, \quad y'(0) = 0.$$

(d) If  $F(s)$  is the complex Fourier transform of  $f(x)$ , then prove that :

$$F\{f(ax)\} = \frac{1}{a} F\left(\frac{s}{a}\right).$$

NA—41—2023

3